

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
12 June 2003 (12.06.2003)

PCT

(10) International Publication Number
WO 03/048387 A2

- (51) International Patent Classification⁷: **C12Q 1/68**
- (21) International Application Number: PCT/GB02/05474
- (22) International Filing Date: 4 December 2002 (04.12.2002)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:
0129012.1 4 December 2001 (04.12.2001) GB
10/227,131 23 August 2002 (23.08.2002) US
- (71) Applicant (for all designated States except US): **SOLEXA LIMITED** [GB/GB]; Chesterford Research Park, Little Chesterford, Nr. Saffron Walden, Essex CB10 1XL (GB).
- (72) Inventors; and
- (75) Inventors/Applicants (for US only): **BALASUBRAMANIAN, Shankar** [GB/GB]; Department of Chemistry, University of Cambridge, Lensfield Road, Cambridge CB2 1EW (GB). **BARNES, Colin** [GB/GB]; Solexa Limited, Chesterford Research Park, Little Chesterford, Nr. Saffron Walden, Essex CB10 1XL (GB). **LIU, Xiaohai** [CN/GB]; Solexa Limited, Chesterford Research Park, Little Chesterford, Nr. Saffron Walden, Essex CB10 1XL (GB). **SWERDLOW, Harold** [US/GB]; Solexa Limited, Chesterford Research Park, Little Chesterford, Nr. Saffron Walden, Essex CB10 1XL (GB).
- (74) Agents: **MURPHY, Colm, Damien et al.**; Boulton Wade Tennant, 70 Gray's Inn Road, London WC1X 8BT (GB).
- (81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.
- (84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).
- Declaration under Rule 4.17:**
— of inventorship (Rule 4.17(iv)) for US only
- Published:**
— without international search report and to be republished upon receipt of that report
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: LABELLED NUCLEOTIDES

(57) Abstract: Nucleosides and nucleotides are disclosed that are linked to detectable labels via a cleavable linker group.

WO 03/048387 A2

LABELLED NUCLEOTIDES

FIELD OF THE INVENTION

5 This invention relates to labelled nucleotides. In particular, this invention discloses nucleotides having a removable label and their use in polynucleotide sequencing methods.

10 BACKGROUND

 Advances in the study of molecules have been led, in part, by improvement in technologies used to characterise the molecules or their biological reactions. In particular, the study of the nucleic acids DNA and RNA has benefited from developing technologies used for sequence analysis and the study of hybridisation events.

 An example of the technologies that have improved the study of nucleic acids, is the development of fabricated arrays of immobilised nucleic acids. These arrays consist typically of a high-density matrix of polynucleotides immobilised onto a solid support material. See, e.g., Fodor et al., *Trends Biotech.* 12:19-26, 1994, which describes ways of assembling the nucleic acids using a chemically sensitized glass surface protected by a mask, but exposed at defined areas to allow attachment of suitably modified nucleotide phosphoramidites. Fabricated arrays can also be manufactured by the technique of "spotting" known polynucleotides onto a solid support at predetermined positions (e.g., Stimpson et al., *Proc. Natl. Acad. Sci. USA* 92:6379-6383, 1995).

 A further development in array technology is the attachment of the polynucleotides to the solid support

- 2 -

material to form single molecule arrays. Arrays of this type are disclosed in International Patent App. WO 00/06770. The advantage of these arrays is that reactions can be monitored at the single molecule level and information on large numbers of single molecules can be collated from a single reaction.

For DNA arrays to be useful, the sequences of the molecules must be determined. U.S. Pat. No. 5,302,509 discloses a method to sequence polynucleotides immobilised on a solid support. The method relies on the incorporation of 3'-blocked bases A, G, C and T having a different fluorescent label to the immobilised polynucleotide, in the presence of DNA polymerase. The polymerase incorporates a base complementary to the target polynucleotide, but is prevented from further addition by the 3'-blocking group. The label of the incorporated base can then be determined and the blocking group removed by chemical cleavage to allow further polymerisation to occur.

Welch et al. (*Chem. Eur. J.* 5(3):951-960, 1999) describes the synthesis of nucleotide triphosphates modified with a 3'-O-blocking group that is photolabile and fluorescent. The modified nucleotides are intended for use in DNA sequencing experiments. However, these nucleotides proved to be difficult to incorporate onto an existing polynucleotide, due to an inability to fit into the polymerase enzyme active site.

Zhu et al. (*Cytometry* 28:206-211, 1997) also discloses the use of fluorescent labels attached to a nucleotide via the base group. The labelled nucleotides are intended for use in fluorescence *in situ* hybridisation (FISH) experiments, where a series of incorporated labelled nucleotides is required to produce a fluorescent "bar code".

- 3 -

SUMMARY OF THE INVENTION

In the present invention, a nucleoside or nucleotide molecule is linked to a detectable label via a cleavable linker group attached to the base, rendering the molecule useful in techniques using labelled nucleosides or nucleotides, e.g., sequencing reactions, polynucleotide synthesis, nucleic acid amplification, nucleic acid hybridization assays, single nucleotide polymorphism studies, and other techniques using enzymes such as polymerases, reverse transcriptases, terminal transferases, or other DNA modifying enzymes. The invention is especially useful in techniques that use labelled dNTPs, such as nick translation, random primer labeling, end-labeling (e.g., with terminal deoxynucleotidyltransferase), reverse transcription, or nucleic acid amplification. The molecules of the present invention are in contrast to the prior art, where the label is attached to the ribose or deoxyribose sugar, or where the label is attached via a non-cleavable linker.

According to a first aspect of the invention, a nucleotide or nucleoside molecule, or an analog thereof, has a base that is linked to a detectable label via a cleavable linker.

The invention features a nucleotide or nucleoside molecule, having a base that is linked to a detectable label via a cleavable linker. The base can be a purine, or a pyrimidine. The base can be a deazapurine. The molecule can have a ribose or deoxyribose sugar moiety. The ribose or deoxyribose sugar can include a protecting group attached via the 2' or 3' oxygen atom. The protecting group can be removed to expose a 3'-OH. The molecule can be a deoxyribonucleotide triphosphate. The detectable label can be a fluorophore. The linker can

- 4 -

be an acid labile linker, a photolabile linker, or can contain a disulphide linkage.

The invention also features a method of labeling a nucleic acid molecule, where the method includes
5 incorporating into the nucleic acid molecule a nucleotide or nucleoside molecule, where the nucleotide or nucleoside molecule has a base that is linked to a detectable label via a cleavable linker. The incorporating step can be accomplished via a terminal
10 transferase, a polymerase or a reverse transcriptase. The base can be a purine, or a pyrimidine. The base can be a deazapurine. The nucleotide or nucleoside molecule can have a ribose or deoxyribose sugar moiety. The ribose or deoxyribose sugar can include a protecting
15 group attached via the 2' or 3' oxygen atom. The protecting group can be removed to expose a 3'-OH group. The molecule can be a deoxyribonucleotide triphosphate. The detectable label can be a fluorophore. The linker can be an acid labile linker, a photolabile linker, or
20 can contain a disulphide linkage. The detectable label and/or the cleavable linker can be of a size sufficient to prevent the incorporation of a second nucleotide or nucleoside into the nucleic acid molecule.

In another aspect, the invention features a method
25 for determining the sequence of a target single-stranded polynucleotide, where the method includes monitoring the sequential incorporation of complementary nucleotides, where the nucleotides each have a base that is linked to a detectable label via a cleavable linker, and where the
30 identity of each nucleotide incorporated is determined by detection of the label linked to the base, and subsequent removal of the label.

The invention also features a method for determining the sequence of a target single-stranded

- 5 -

polynucleotide, where the method includes: (a) providing nucleotides, where the nucleotides have a base that is linked to a detectable label via a cleavable linker, and where the detectable label linked to each type of nucleotide can be distinguished upon detection from the detectable label used for other types of nucleotides; (b) incorporating a nucleotide into the complement of the target single stranded polynucleotide; (c) detecting the label of the nucleotide of (b), thereby determining the type of nucleotide incorporated; (d) removing the label of the nucleotide of (b); and (e) optionally repeating steps (b) - (d) one or more times; thereby determining the sequence of a target single-stranded polynucleotide.

In the methods described herein, each of the nucleotides can be brought into contact with the target sequentially, with removal of non-incorporated nucleotides prior to addition of the next nucleotide, where detection and removal of the label is carried out either after addition of each nucleotide, or after addition of all four nucleotides.

In the methods, all of the nucleotides can be brought into contact with the target simultaneously, i.e., a composition comprising all of the different nucleotides is brought into contact with the target, and non-incorporated nucleotides are removed prior to detection and subsequent to removal of the label(s).

The methods can comprise a first step and a second step, where in the first step, a first composition comprising two of the four nucleotides is brought into contact with the target, and non-incorporated nucleotides are removed prior to detection and subsequent to removal of the label, and where in the second step, a second composition comprising the two

- 6 -

nucleotides not included in the first composition is brought into contact with the target, and non-incorporated nucleotides are removed prior to detection and subsequent to removal of the label, and where the first steps and the second step can be optionally repeated one or more times.

The methods described herein can also comprise a first step and a second step, where in the first step, a composition comprising one of the four nucleotides is brought into contact with the target, and non-incorporated nucleotides are removed prior to detection and subsequent to removal of the label, and where in the second step, a second composition comprising the three nucleotides not included in the first composition is brought into contact with the target, and non-incorporated nucleotides are removed prior to detection and subsequent to removal of the label, and where the first steps and the second step can be optionally repeated one or more times.

The methods described herein can also comprise a first step and a second step, where in the first step, a first composition comprising three of the four nucleotides is brought into contact with the target, and non-incorporated nucleotides are removed prior to detection and subsequent to removal of the label, and where in the second step, a composition comprising the nucleotide not included in the first composition is brought into contact with the target, and non-incorporated nucleotides are removed prior to detection and subsequent to removal of the label, and where the first steps and the second step can be optionally repeated one or more times.

In a further aspect, the invention features a kit, where the kit includes: (a) individual the nucleotides,

- 7 -

where each nucleotide has a base that is linked to a detectable label via a cleavable linker, and where the detectable label linked to each nucleotide can be distinguished upon detection from the detectable label used for other three nucleotides; and (b) packaging materials therefor. The kit can further include an enzyme and buffers appropriate for the action of the enzyme.

The nucleotides/nucleosides are suitable for use in many different DNA-based methodologies, including DNA synthesis and DNA sequencing protocols.

According to another aspect of the invention, a method for determining the sequence of a target polynucleotide comprises monitoring the sequential incorporation of complementary nucleotides, wherein the nucleotides comprise a detectable label linked to the base portion of the nucleotide via a cleavable linker, incorporation is detected by monitoring the label, and the label is removed to permit further nucleotide incorporation to occur.

DESCRIPTION OF THE DRAWINGS

Fig. 1 shows exemplary nucleotide structures useful in the invention. For each structure, X can be H, phosphate, diphosphate or triphosphate. R_1 and R_2 can be the same or different, and can be selected from H, OH, or any group which can be transformed into an OH.

Fig. 2 shows structures of linkers useful in the invention, including (1) disulfide linkers and acid labile linkers, (2) dialkoxybenzyl linkers, (3) Sieber linkers, (4) indole linkers and (5) t-butyl Sieber linkers in addition to a general definition of the linkers that may be used.

- 8 -

Fig. 3 shows some functional molecules useful in the invention, including some cleavable linkers. In these structures, R_1 and R_2 may be the same or different, and can be H, OH, or any group which can be transformed into an OH group, including a carbonyl. R_3 represents one or more substituents independently selected from alkyl, alkoxy, amino or halogen groups. Alternatively, cleavable linkers may be constructed from any labile functionality used on the 3'-block.

Fig. 4 shows a denaturing gel showing the incorporation of the triphosphate of Example 1 using Klenow polymerase.

Fig. 5 shows a denaturing gel showing the incorporation of the triphosphate of Example 3 using Klenow polymerase.

Fig. 6 shows a denaturing gel showing the incorporation of the triphosphate of Example 4 using Klenow polymerase.

DETAILED DESCRIPTION

The present invention relates to nucleotides and nucleosides that are modified by attachment of a label via a cleavable linker, thereby rendering the molecule useful in techniques where the labelled molecule is to interact with an enzyme, such as sequencing reactions, polynucleotide synthesis, nucleic acid amplification, nucleic acid hybridization assays, single nucleotide polymorphism studies, techniques using enzymes such as polymerase, reverse transcriptase, terminal transferase, techniques that use labelled dNTPs (e.g., nick translation, random primer labeling, end-labeling (e.g., with terminal deoxynucleotidyltransferase), reverse transcription, or nucleic acid amplification).

- 9 -

As is known in the art, a "nucleotide" consists of a nitrogenous base, a sugar, and one or more phosphate groups. In RNA, the sugar is a ribose, and in DNA is a deoxyribose, *i.e.*, a sugar lacking a hydroxyl group that is present in ribose. The nitrogenous base is a derivative of purine or pyrimidine. The purines are adenosine (A) and guanine (G), and the pyrimidines are cytosine (C) and thymine (T) (or in the context of RNA, uracil (U)). The C-1 atom of deoxyribose is bonded to N-1 of a pyrimidine or N-9 of a purine. A nucleotide is also a phosphate ester of a nucleoside, with esterification occurring on the hydroxyl group attached to C-5 of the sugar. Nucleotides are usually mono, di- or triphosphates.

A "nucleoside" is structurally similar to a nucleotide, but is missing the phosphate moieties. An example of a nucleoside analog would be one in which the label is linked to the base and there is no phosphate group attached to the sugar molecule.

Although the base is usually referred to as a purine or pyrimidine, the skilled person will appreciate that derivatives and analogs are available which do not alter the capability of the nucleotide or nucleoside to undergo Watson-Crick base pairing. "Derivative" or "analog" means a compound or molecule whose core structure is the same as, or closely resembles that of, a parent compound, but which has a chemical or physical modification, such as a different or additional side groups, which allows the derivative nucleotide or nucleoside to be linked to another molecule. For example, the base can be a deazapurine. The derivatives should be capable of undergoing Watson-Crick pairing. "Derivative" and "analog" also mean a synthetic nucleotide or nucleoside derivative having modified base

- 10 -

moieties and/or modified sugar moieties. Such derivatives and analogs are discussed in, e.g., Scheit, *Nucleotide Analogs* (John Wiley & Son, 1980) and Uhlman et al., *Chemical Reviews* 90:543-584, 1990. Nucleotide
5 analogs can also comprise modified phosphodiester linkages, including phosphorothioate, phosphorodithioate, alkylphosphonate, phosphoranilidate and phosphoramidate linkages. The analogs should be capable of undergoing Watson-Crick base pairing.
10 "Derivative" and "analog", as used herein, may be used interchangeably, and are encompassed by the terms "nucleotide" and "nucleoside" as defined herein.

The present invention can make use of conventional detectable labels. Detection can be carried out by any
15 suitable method, including fluorescence spectroscopy or by other optical means. The preferred label is a fluorophore, which, after absorption of energy, emits radiation at a defined wavelength. Many suitable fluorescent labels are known. For example, Welch et al.
20 (*Chem. Eur. J.* 5(3):951-960, 1999) discloses dansyl-functionalised fluorescent moieties that can be used in the present invention. Zhu et al. (*Cytometry* 28:206-211, 1997) describes the use of the fluorescent labels Cy3 and Cy5, which can also be used in the present
25 invention. Labels suitable for use are also disclosed in Prober et al. (*Science* 238:336-341, 1987); Connell et al. (*BioTechniques* 5(4):342-384, 1987), Ansorge et al. (*Nucl. Acids Res.* 15(11):4593-4602, 1987) and Smith et al. (*Nature* 321:674, 1986). Other commercially
30 available fluorescent labels include, but are not limited to, fluorescein, rhodamine (including TMR, texas red and Rox), alexa, bodipy, acridine, coumarin, pyrene, benzanthracene and the cyanins.

- 11 -

Multiple labels can also be used in the invention. For example, bi-fluorophore FRET cassettes (*Tet. Letts.* 46:8867-8871, 2000) are well known in the art and can be utilised in the present invention. Multi-fluor
5 dendrimeric systems (*J. Amer. Chem. Soc.* 123:8101-8108, 2001) can also be used.

Although fluorescent labels are preferred, other forms of detectable labels will be apparent as useful to those of ordinary skill. For example, microparticles,
10 including quantum dots (Empodocles, *et al.*, *Nature* 399:126-130, 1999), gold nanoparticles (Reichert *et al.*, *Anal. Chem.* 72:6025-6029, 2000), microbeads (Lacoste *et al.*, *Proc. Natl. Acad. Sci USA* 97(17):9461-9466, 2000), and tags detectable by mass spectrometry can all be
15 used.

Multi-component labels can also be used in the invention. A multi-component label is one which is dependent on the interaction with a further compound for detection. The most common multi-component label used
20 in biology is the biotin-streptavidin system. Biotin is used as the label attached to the nucleotide base. Streptavidin is then added separately to enable detection to occur. Other multi-component systems are available. For example, dinitrophenol has a
25 commercially available fluorescent antibody that can be used for detection.

The label (or label and linker construct) can be of a size or structure sufficient to act as a block to the incorporation of a further nucleotide onto the
30 nucleotide of the invention. This permits controlled polymerization to be carried out. The block can be due to steric hindrance, or can be due to a combination of size, charge and structure.

The invention will be further described with

- 12 -

reference to nucleotides. However, unless indicated otherwise, the reference to nucleotides is also intended to be applicable to nucleosides. The invention will also be further described with reference to DNA, although the description will also be applicable to RNA, PNA, and other nucleic acids, unless otherwise indicated.

The modified nucleotides of the invention use a cleavable linker to attach the label to the nucleotide. The use of a cleavable linker ensures that the label can, if required, be removed after detection, avoiding any interfering signal with any labelled nucleotide incorporated subsequently.

Cleavable linkers are known in the art, and conventional chemistry can be applied to attach a linker to a nucleotide base and a label. The linker can be cleaved by any suitable method, including exposure to acids, bases, nucleophiles, electrophiles, radicals, metals, reducing or oxidising agents, light, temperature, enzymes etc. Suitable linkers can be adapted from standard chemical blocking groups, as disclosed in Greene & Wuts, *Protective Groups in Organic Synthesis*, John Wiley & Sons. Further suitable cleavable linkers used in solid-phase synthesis are disclosed in Guillier et al. (*Chem. Rev.* 100:2092-2157, 2000).

The use of the term "cleavable linker" is not meant to imply that the whole linker is required to be removed from the nucleotide base. The cleavage site can be located at a position on the linker that ensures that part of the linker remains attached to the nucleotide base after cleavage.

The linker can be attached at any position on the nucleotide base provided that Watson-Crick base pairing

- 13 -

can still be carried out. In the context of purine bases, it is preferred if the linker is attached via the 7 position of the purine or the preferred deazapurine analogue, via an 8-modified purine, via an N-6 modified adenosine or an N-2 modified guanine. For pyrimidines, attachment is preferably via the 5 position on cytidine, thymidine or uracil and the N-4 position on cytosine. Suitable nucleotide structures are shown in Fig. 1. For each structure in Fig. 1, X can be H, phosphate, diphosphate or triphosphate. R_1 and R_2 can be the same or different, and can be selected from H, OH, or any group which can be transformed into an OH, including, but not limited to, a carbonyl.

Suitable linkers are shown generally in Fig. 2 and include, but are not limited to, disulfide linkers (1), acid labile linkers (2, 3, 4 and 5; including dialkoxybenzyl linkers (e.g., 2), Sieber linkers (e.g., 3), indole linkers (e.g., 4), *t*-butyl Sieber linkers (e.g., 5)), electrophilically cleavable linkers, nucleophilically cleavable linkers, photocleavable linkers, cleavage under reductive conditions, oxidative conditions, cleavage via use of safety-catch linkers, and cleavage by elimination mechanisms.

25 A. Electrophilically cleaved linkers.

Electrophilically cleaved linkers are typically cleaved by protons and include cleavages sensitive to acids. Suitable linkers include the modified benzylic systems such as trityl, *p*-alkoxybenzyl esters and *p*-alkoxybenzyl amides. Other suitable linkers include *tert*-butyloxycarbonyl (Boc) groups and the acetal system (e.g., as is shown in Fig. 3 as $O-C(R_4)(R_5)-O-R_6$).

The use of thiophilic metals, such as nickel, silver or mercury, in the cleavage of thioacetal or

- 14 -

other sulphur-containing protecting groups can also be considered for the preparation of suitable linker molecules.

5 B. Nucleophilically cleaved linkers.

 Nucleophilic cleavage is also a well recognised method in the preparation of linker molecules. Groups such as esters that are labile in water (i.e., can be cleaved simply at basic pH) and groups that are labile to non-aqueous nucleophiles, can be used. Fluoride ions can be used to cleave silicon-oxygen bonds in groups such as triisopropyl silane (TIPS) or t-butyldimethyl silane (TBDMS).

15 C. Photocleavable linkers.

 Photocleavable linkers have been used widely in carbohydrate chemistry. It is preferable that the light required to activate cleavage does not affect the other components of the modified nucleotides. For example, if a fluorophore is used as the label, it is preferable if this absorbs light of a different wavelength to that required to cleave the linker molecule. Suitable linkers include those based on O-nitrobenzyl compounds and nitroveratryl compounds. Linkers based on benzoin chemistry can also be used (Lee et al., *J. Org. Chem.* 64:3454-3460, 1999).

 D. Cleavage under reductive conditions

 There are many linkers known that are susceptible to reductive cleavage. Catalytic hydrogenation using palladium-based catalysts has been used to cleave benzyl and benzyloxycarbonyl groups. Disulphide bond reduction is also known in the art.

- 15 -

E. Cleavage under oxidative conditions

Oxidation-based approaches are well known in the art. These include oxidation of *p*-alkoxybenzyl groups and the oxidation of sulphur and selenium linkers. The use of aqueous iodine to cleave disulphides and other sulphur or selenium-based linkers is also within the scope of the invention.

F. Safety-catch linkers

Safety-catch linkers are those that cleave in two steps. In a preferred system the first step is the generation of a reactive nucleophilic center followed by a second step involving an intra-molecular cyclization that results in cleavage. For example, levulinic ester linkages can be treated with hydrazine or photochemistry to release an active amine, which can then be cyclised to cleave an ester elsewhere in the molecule (Burgess et al., *J. Org. Chem.* 62:5165-5168, 1997).

G. Cleavage by elimination mechanisms

Elimination reactions can also be used. For example, the base-catalysed elimination of groups such as Fmoc and cyanoethyl, and palladium-catalysed reductive elimination of allylic systems, can be used.

As well as the cleavage site, the linker can comprise a spacer unit. The spacer distances the nucleotide base from the cleavage site or label. The length of the linker is unimportant provided that the label is held a sufficient distance from the nucleotide so as not to interfere with any interaction between the nucleotide and an enzyme.

The modified nucleotides can also comprise additional groups or modifications to the sugar group. For example, a dideoxyribose derivative, lacking two

- 16 -

oxygen on the ribose ring structure (at the 2' and 3' positions), can be prepared and used as a block to further nucleotide incorporation on a growing oligonucleotide strand. The protecting group is intended to prevent nucleotide incorporation onto a nascent polynucleotide strand, and can be removed under defined conditions to allow polymerisation to occur. In contrast to the prior art, there is no detectable label attached at the ribose 3' position. This ensures that steric hindrance with the polymerase enzyme is reduced, while still allowing control of incorporation using the protecting group.

The skilled person will appreciate how to attach a suitable protecting group to the ribose ring to block interactions with the 3'-OH. The protecting group can be attached directly at the 3' position, or can be attached at the 2' position (the protecting group being of sufficient size or charge to block interactions at the 3' position). Alternatively, the protecting group can be attached at both the 3' and 2' positions, and can be cleaved to expose the 3'OH group.

Suitable protecting groups will be apparent to the skilled person, and can be formed from any suitable protecting group disclosed in Green and Wuts, *supra*. The protecting group should be removable (or modifiable) to produce a 3' OH group. The process used to obtain the 3' OH group can be any suitable chemical or enzymic reaction.

The labile linker may consist of functionality cleavable under identical conditions to the block. This will make the deprotection process more efficient as only a single treatment will be required to cleave both the label and the block. Thus the linker may contain functional groups as described in Fig. 3, which could be

- 17 -

cleaved with the hydroxyl functionality on either the residual nucleoside or the removed label. The linker may also consist of entirely different chemical functionality that happens to be labile to the conditions used to cleave the block.

The term "alkyl" covers both straight chain and branched chain alkyl groups. Unless the context indicates otherwise, the term "alkyl" refers to groups having 1 to 8 carbon atoms, and typically from 1 to 6 carbon atoms, for example from 1 to 4 carbon atoms. Examples of alkyl groups include methyl, ethyl, propyl, isopropyl, n-butyl, isobutyl, tert-butyl, n-pentyl, 2-pentyl, 3-pentyl, 2-methyl butyl, 3-methyl butyl, and n-hexyl and its isomers.

Examples of cycloalkyl groups are those having from 3 to 10 ring atoms, particular examples including those derived from cyclopropane, cyclobutane, cyclopentane, cyclohexane and cycloheptane, bicycloheptane and decalin.

Examples of alkenyl groups include, but are not limited to, ethenyl (vinyl), 1-propenyl, 2-propenyl (allyl), isopropenyl, butenyl, buta-1,4-dienyl, pentenyl, and hexenyl.

Examples of cycloalkenyl groups include, but are not limited to, cyclopropenyl, cyclobutenyl, cyclopentenyl, cyclopentadienyl and cyclohexenyl.

The term alkoxy refers to C₁₋₆ alkoxy unless otherwise indicated: -OR, wherein R is a C₁₋₆alkyl group. Examples of C₁₋₆ alkoxy groups include, but are not limited to, -OMe (methoxy), -OEt (ethoxy), -O(nPr) (n-propoxy), -O(iPr) (isopropoxy), -O(nBu) (n-butoxy), -O(sBu) (sec-butoxy), -O(iBu) (isobutoxy), and -O(tBu) (tert-butoxy).

- 18 -

The term amino refers to groups of type NR^1R^2 , wherein R^1 and R^2 are independently selected from hydrogen, a C_{1-6} alkyl group (also referred to as C_{1-6} alkylamino or di- C_{1-6} alkylamino).

5 The term "halogen" as used herein includes fluorine, chlorine, bromine and iodine.

The nucleotide molecules of the present invention are suitable for use in many different methods where the detection of nucleotides is required.

10 DNA sequencing methods, such as those outlined in U.S. Pat. No. 5,302,509 can be carried out using the nucleotides.

15 A method for determining the sequence of a target polynucleotide can be carried out by contacting the target polynucleotide separately with the different nucleotides to form the complement to that of the target polynucleotide, and detecting the incorporation of the nucleotides. Such a method makes use of polymerisation, whereby a polymerase enzyme extends the complementary strand by incorporating the correct nucleotide complementary to that on the target. The polymerisation reaction also requires a specific primer to initiate polymerisation.

20 For each cycle, the incorporation of the labelled nucleotide is carried out by the polymerase enzyme, and the incorporation event is then determined. Many different polymerase enzymes exist, and it will be evident to the person of ordinary skill which is most appropriate to use. Preferred enzymes include DNA
25 polymerase I, the Klenow fragment, DNA polymerase III, T4 or T7 DNA polymerase, Taq polymerase or vent polymerase. A polymerase engineered to have specific properties can also be used.

The sequencing methods are preferably carried out

- 19 -

with the target polynucleotide arrayed on a solid support. Multiple target polynucleotides can be immobilised on the solid support through linker molecules, or can be attached to particles, e.g.,
5 microspheres, which can also be attached to a solid support material.

The polynucleotides can be attached to the solid support by a number of means, including the use of biotin-avidin interactions. Methods for immobilizing
10 polynucleotides on a solid support are well known in the art, and include lithographic techniques and "spotting" individual polynucleotides in defined positions on a solid support. Suitable solid supports are known in the art, and include glass slides and beads, ceramic and
15 silicon surfaces and plastic materials. The support is usually a flat surface although microscopic beads (microspheres) can also be used and can in turn be attached to another solid support by known means. The microspheres can be of any suitable size, typically in
20 the range of from 10 nm to 100 nm in diameter. In a preferred embodiment, the polynucleotides are attached directly onto a planar surface, preferably a planar glass surface. Attachment will preferably be by means of a covalent linkage. Preferably, the arrays that are
25 used are single molecule arrays that comprise polynucleotides in distinct optically resolvable areas, e.g., as disclosed in International App. No. WO 00/06770.

The sequencing method can be carried out on both
30 single polynucleotide molecule and multi-polynucleotide molecule arrays, i.e., arrays of distinct individual polynucleotide molecules and arrays of distinct regions comprising multiple copies of one individual polynucleotide molecule. Single molecule arrays allow

- 20 -

each individual polynucleotide to be resolved separately. The use of single molecule arrays is preferred. Sequencing single molecule arrays non-destructively allows a spatially addressable array to be
5 formed.

The method makes use of the polymerisation reaction to generate the complementary sequence of the target. The conditions necessary for polymerisation to occur will be apparent to the skilled person.

10 To carry out the polymerase reaction it will usually be necessary to first anneal a primer sequence to the target polynucleotide, the primer sequence being recognised by the polymerase enzyme and acting as an initiation site for the subsequent extension of the
15 complementary strand. The primer sequence may be added as a separate component with respect to the target polynucleotide. Alternatively, the primer and the target polynucleotide may each be part of one single stranded molecule, with the primer portion forming an
20 intramolecular duplex with a part of the target, i.e., a hairpin loop structure. This structure may be immobilised to the solid support at any point on the molecule. Other conditions necessary for carrying out the polymerase reaction, including temperature, pH,
25 buffer compositions etc., will be apparent to those skilled in the art.

The modified nucleotides of the invention are then brought into contact with the target polynucleotide, to allow polymerisation to occur. The nucleotides may be
30 added sequentially, i.e., separate addition of each nucleotide type (A, T, G or C), or added together. If they are added together, it is preferable for each nucleotide type to be labelled with a different label.

- 21 -

This polymerisation step is allowed to proceed for a time sufficient to allow incorporation of a nucleotide.

5 Nucleotides that are not incorporated are then removed, for example, by subjecting the array to a washing step, and detection of the incorporated labels may then be carried out.

10 Detection may be by conventional means, for example if the label is a fluorescent moiety, detection of an incorporated base may be carried out by using a confocal scanning microscope to scan the surface of the array with a laser, to image a fluorophore bound directly to the incorporated base. Alternatively, a sensitive 2-D detector, such as a charge-coupled detector (CCD), can be used to visualise the individual signals generated. 15 However, other techniques such as scanning near-field optical microscopy (SNOM) are available and may be used when imaging dense arrays. For example, using SNOM, individual polynucleotides may be distinguished when separated by a distance of less than 100 nm, e.g., 10 nm to 10 μ m. For a description of scanning near-field optical microscopy, see Moyer et al., *Laser Focus World* 29:10, 1993. Suitable apparatus used for imaging polynucleotide arrays are known and the technical set-up 20 will be apparent to the skilled person. 25

After detection, the label may be removed using suitable conditions that cleave the linker.

30 The use of the modified nucleotides is not limited to DNA sequencing techniques, and other techniques, including polynucleotide synthesis, DNA hybridisation assays and single nucleotide polymorphism studies, may also be carried out using nucleotides of the invention. Any technique that involves the interaction between a nucleotide and an enzyme may make use of the molecules


- 22 -

of the invention. For example, the molecule may be used as a substrate for a reverse transcriptase or terminal transferase enzyme.


Suitable structures are described in the following
 5 Examples and are shown in the accompanying drawings.

EXAMPLES


Example 1. Synthesis of Disulfide Linker.



- 23 -

**1a**

tButyl-N-(2-mercaptoethyl) carbamate (3 mmol, 0.5 mL) was added dropwise to a solution of 1.32g (6.0 mmol) aldrithiol in 15 mL MeOH. After 1.5 h the reaction had gone to completion and the solvent was evaporated. The crude product was purified by chromatography on silica with ethyl acetate:petroleum ether (1:4). Product **1a** was obtained as a slightly yellow oil (0.76 g, 2.67 mmol, 89 %). ¹H NMR (500 Mhz, D₆-DMSO): δ=1.38 (s, 9 H, tBu), 2.88 (t, J = 6.6 Hz, 2 H, SCH₂) 3.20 (q, J = 6.6 Hz, 2 H, CH₂NH), 7.02 (bs, 1 H, NH), 7.24 (ddd, J = 7.3 Hz, J = 4.9 Hz, J = 1.0 Hz, 1 H, H-5), 7.77 (dt, J = 8.1 Hz, J = 1.0 Hz, 1 H, H-3), 7.82 (ddd, J = 8.1 Hz, J = 7.4 Hz, J = 1.8 Hz, 1 H, H-4), 8.46 (ddd, J = 4.9 Hz, J = 1.8 Hz, J = 1.0 Hz, 1 H, H-6).


**1b**

- 24 -

To deprotect the amine of **1a**, 17 mg of **1a** (60 μ mol) was dissolved in a mixture of 0.5 mL DCM and 0.5 mL trifluoroacetic acid. This mixture was stirred for 2.5h at room temperature and then the solvents were removed under reduced pressure. The residue was three times redissolved in 2 mL DCM and evaporated to dryness. The deprotected product was dried under high vacuum for 3 h and then dissolved in 1 mL dry DMF. It was assumed that the deprotection had gone to completion.

To a solution of 15 mg 5-carboxy tetra methyl rhodamine (35 μ mol) in 2 mL DMF were added 8.0 mg N-hydroxy succinimide (70 μ mol) and 7.8 mg DCC (38 μ mol). The mixture was stirred for 6 h in the dark. Then 22 μ l DIPEA (126 μ mol) and the solution of deprotected **1a** in 1 mL DMF were added. After stirring the reaction mixture overnight in the dark, the solvent was removed under reduced pressure. The residue was dissolved in DCM and washed with saturated NaCl solution. After drying over MgSO_4 , the crude mixture was purified on silica with CHCl_3 :MeOH (3:1) as solvent. **1b** was isolated as a dark red solid in 90 % yield (19.2 mg, 31.4 μ mol). ^1H NMR (500 MHz, D_6 -DMSO): δ = 3.09 (t, J = 6.7 Hz, 2 H, SCH_2), 3.63 (q, J = 6.2 Hz, 2 H, CH_2NH), 6.48 - 6.53 (m, 6 H, H-Anthracene), 7.23-7.26 [m, 1 H, H-5 (pyridine)], 7.32 (d, J = 7.9 Hz, 1 H, H-3), 7.81 - 7.82 [m, 2 H, H-3 +H-4 (pyridine)], 8.21 (d, J = 7.9 Hz, 1 H, H-4), 8.43 (s, 1 H, H-6), 8.47 [dt, J = 4.7 Hz, J = 1.3 Hz, 1 H, H-6 (pyridine)], 9.03 (t, J = 5.2 Hz, 1 H, NH).


- 25 -



Mercaptopropionic acid (20.6 μ mol, 1.8 ml) was added
 to a solution of 19.6 mg **1b** (32.7 μ mol) in 2 mL MeOH.
 5 The mixture was stirred for 2.5 h in the dark. The
 solvent was removed under reduced pressure. The crude
 product was purified by chromatography on silica with
 CHCl₃:MeOH:AcOH 15:1:0.5 as the solvent mixture.
 15.5 mg (26 μ mol, 80 %) dark red crystals **1c** could be
 10 isolated. ¹H NMR (500 MHz, D₂O): δ = 2.53 (t,
 J = 7.0 Hz, 2 H, CH₂COOH), 2.88 (t, J = 7.0 Hz, 2 H,
 CH₂CH₂COOH), 2.96 - 2.99 (m, 2 H, CH₂CH₂NH), 3.73 (t,
 J = 6.3 Hz, 2 H, CH₂NH), 6.53 (d, J = 2.4 Hz, 2 H, H-
 Anthracene), 6.81 (dd, J = 9.5 Hz, J = 4.5 Hz, 2 H, H-
 15 Anthracene), 7.12 (d, J = 9.5 Hz, 2 H, H-Anthracene),
 7.48 (d, J = 7.9 Hz, 1 H, H-3), 7.95 (dd, J = 8.1 Hz,
 J = 1.9 Hz, 1 H, H-2) 8.13 (d, J = 1.9 Hz, 1 H, H-1).
 +ve electro spray (C₃₀H₃₁N₃O₆S₂): expected 593.17; found
 594.3 [M+H], 616.2 [M+Na].

20

- 26 -



To a solution of 25.8 mg **1c** (43.4 μmol) in 3 mL DMF (dry) were added 9.9 mg N-hydroxy succinimide (86.8 μmol) and 9.7 mg DCC (47.1 μmol). The mixture was stirred in the dark for 5 h at room temperature and then put in the fridge overnight. The mixture was filtered through a plug of cotton wool in a new flask and to this was added a solution of 865 μl propargylamino dUTP (14.7 μmol , 17 μmol in 1 mL H_2O) and 3 mL sodium borate buffer (0.1 M solution, pH 9). The mixture was stirred overnight. After removal of solvents the residue was dissolved in as little water as possible and purified by HPLC. A Zorbax C18 column was used with 0.1 M triethyl ammonium bicarbonate (TEAB) and acetonitrile as buffers.

^{31}P NMR (400 MHz, D_2O): δ = -4.73 (d), -9.93 (d), 19.03 (t). -ve electro spray ($\text{C}_{42}\text{H}_{47}\text{N}_6\text{O}_{19}\text{P}_3\text{S}_2$ assuming 4 H^+ counter ions): expected 1096.16; found 1092.9. UV in Water: $\lambda_{(\text{max})}$ = 555 nm $A_{(555)}$ = 0.885 (c = 0.036 μmol).

20


Triphosphate (**1**) was successfully incorporated

- 27 -

using Klenow DNA polymerase. The reaction was performed in the following conditions: 50 mM Tris.HCl (pH 7.5), 10 mM NaCl, 2 mM DTT, 0.1 mM EDTA, 5 mM MgCl₂, 2 μM compound 3, 100 nM DNA template (previously labelled with P32 and T4 polynucleotide kinase) and 10 units of commercial exo-Klenow (Amersham Corp., Arlington Heights, Illinois, USA). The DNA templates were self-complementary hairpins (5' - TACCgTCgACgTCgACgCTggCg-AgCgTgCTgCggTTTTT(C6-amino)TTACCgCAGCACgCTCgCCAgCg; SEQ ID NO:1). The reaction was performed in 100 μL volume at 37 °C with timepoints taken at 0, 1, 3, 5 and 10 min. The reaction products were electrophoresed down a denaturing (8 M urea) 20 % polyacrylamide gel and imaged on a typhoon phosphorimager. Complete single base extension was seen in 1 minute indicating efficient polymerase incorporation (disulfide linker gel, Fig. 4). A second set of lanes is shown in which the material is exposed to DTT after the incorporation. A different band shift can be seen which shows removal of the dye from the DNA construct, thus a cycle of polymerase incorporation and cleavage has been shown using this disulfide compound.

- 28 -


Example 2. Synthesis of TMR-Sieber Linker Free Acid.



5 - [- 9 - [9 - (f l u o r e n y l -
 5 methyloxycarbonyl)amino]xanthene-3-yl]valeric acid, (42.8
 mg, 80 μ mol) was stirred at room temperature with
 disuccinimidyl carbonate (22.5 mg, 88 μ mol) and *N,N*-
 dimethyl aminopyridine (10.8 mg, 88 μ mol) in DMF. After
 5 minutes, mono-5-carboxy TMR ethylene diamine (198.9
 10 mg, 40 μ mol) was added followed by DIPEA (13.9 μ l, 80
 μ mol). The reaction was stirred at room temperature.
 After 2 hrs, the reaction mixture was diluted with
 dichloromethane (100 mL) and the resulting solution was
 extracted with 1 M aqueous potassium dihydrogen
 15 phosphate (50 mL). The DCM layer was separated and
 evaporated under reduced pressure. The residue was
 purified by a short column chromatography. The
 fractions eluting with 40 % methanol in chloroform were
 collected and evaporated under reduced pressure. The
 20 residue was then dissolved in dry DMF (1 mL) and *N*-(2-
 mercaptoethyl)aminomethyl polystyrene (200 mg, 400 μ mol)
 and DBU (12 μ l, 80 μ mol). After 10 minutes at room
 temperature, the resins were filtered off and rinsed
 with dry DMF (1 mL). All the filtrates were combined
 25 and then added to a solution of succinic anhydride (80

mg, 800 μ mol), DIPEA (139 μ l, 800 μ mol) and DMAP (9.8 mg, 80 μ mol) in DMF (1 mL). The reaction mixture was then stirred at room temperature. After overnight (16 hrs), all the solvents were evaporated under reduced pressure and the residue was purified by a short column chromatography. The title compound eluted with 30 % methanol in chloroform obtained as purple powders (22 mg, overall yield 63 %). ^1H NMR [D_6 -DMSO]: 8.82 (1H, t, J 5.4, ex.), 8.75 (1H, d, J 8.9, ex.), 8.42 (1H, d, J 1.5), 8.20 (1H, dd, J 8.0 and 1.5), 7.95 (1H, t, J 5.9, ex.), 7.34 (1H, d, J 7.3), 7.30-7.27 (2H, m), 7.21 (1H, d, J 8.5), 7.16-7.07 (2H, m), 6.68 (1H, dd, J 8.8 and 2.5), 6.65 (1H, d, J 2.4), 6.49-6.43 (6H, m), 6.18 (1H, d, J 5.6), 3.95 (1H, t, J 5.9), 3.39-3.36 (2H, m), 3.30-3.27 (2H, m), 2.92 (12H, s), 2.37-2.33 (2H, m), 2.14 (2H, t, J 7.2) and 1.70-1.62 (4H, m). MS[(ES(+))], m/z 868.5 (MH^+).

Example 3. Synthesis of TMR-Sieber Linker-dUTP(3)



- 30 -

TMR-sieber linker free acid (4.34 mg, 5 μ mol) was stirred with disuccinimidyl carbonate (1.74 mg, 7.5 μ mol) and *N,N*-dimethyl aminopyridine (0.92 mg, 7.5 μ mol) in DMF (1 mL) at room temperature. After 10 minutes, all the reaction mixture was added to tetra-(tributylammonium) salt of 5-(3-aminopropynyl)-2'-deoxyuridine-5'-triphosphate (10 μ mol). The reaction was stirred at room temperature for 4 hrs and stored in the fridge overnight. The reaction mixture was then diluted with chilled water (10 mL) and all the resulting solution was applied onto a short column of DEAE A-25. The column was initially eluted with 0.1 M TEAB buffer and then 0.7 M TEAB buffer. The 0.7 M TEAB eluents were collected and evaporated under reduced pressure. The residue was co-evaporated with MeOH (2 x 10 mL) and then purified by preparative HPLC. The title compound was obtained as triethylammonium salt in 31 % yield (based on the quantification of TMR at 555 nm in water (pH 7)). ¹HNMR in D₂O indicated two diastereoisomers, due to the sieber linker moiety and there were approximately three triethylammonium count ions. ¹HNMR [D₂O]: 8.18 (1H, m), 8.06 (1H, m), 7.76 (0.55H, s), 7.74 (0.45H, s), 7.36-7.09 (5H, m), 6.89-6.72 (3H, m), 6.59-6.37 (5H, m), 6.12 (0.55H, t, *J* 6.6), 6.05 (0.45H, t, *J* 6.6), 5.99 (0.45H, d, *J* 2.5), 5.91 (1.1H, m), 5.88 (0.45H, s), 4.49 (0.55H, m), 4.43 (0.45H, m), 4.00-3.35 (9H, m), 3.30-2.95 (32H, m), 2.65-2.52 (4H, m), 2.25-2.05 (4H, m), 1.62-1.42 (4H, m) and 1.23 (27H, t, *J* 7.3). ³¹P [D₂O]: -9.91 (^νP, d, *J* 19.2), [-11.08 (^αP, d, *J* 20.1) and -11.30 (^αP, d, *J* 20.1), due to two diastereoisomers] and -22.57 (^βP, m). MS[(ES(-))], *m/z* 1369.1 (M⁻).


Triphosphate (3) was successfully incorporated using Klenow DNA polymerase. The reaction was performed in the following conditions: 50 mM Tris.HCl (pH 7.5), 10

- 31 -

mM NaCl, 2 mM DTT, 0.1 mM EDTA, 5 mM MgCl₂, 2 μM compound
 3, 100 nM DNA template (previously labelled with P32 and
 T4 polynucleotide kinase) and 10 units of commercial
 5 exo- Klenow (Amersham Corp. Arlington Heights, Illinois,
 USA). The DNA templates were self-complementary
 hairpins (5' - TACCgTCgACgTCgACgCTggCg -
 AgCgTgCTgCggTTTTT(C6-amino)TTACCgCAgCACgCTCgCCAgCg; SEQ
 ID NO:1). The reaction was performed in 100 μL volume
 at 37 °C with timepoints taken at 0, 1, 3, 5 and 10 min.
 10 The reaction products were electrophoresed down a
 denaturing (8 M urea) 20 % polyacrylamide gel and imaged
 on a typhoon phosphorimager. Complete single base
 extension was seen in 1 minute indicating efficient
 polymerase incorporation (Sieber linker gel, Fig. 5).

15

Example 4. Synthesis of TMR-Indole Linker-dUTP (4)



20 Triphosphate (4) was successfully incorporated
 using Klenow DNA polymerase. The reaction was performed
 in the following conditions: 50 mM Tris.HCl (pH 7.5), 10
 mM NaCl, 2 mM DTT, 0.1 mM EDTA, 5 mM MgCl₂, 2 μM compound

- 32 -

3, 100 nM DNA template (previously labelled with P32 and T4 polynucleotide kinase) and 10 units of commercial exo-Klenow (Amersham Corp., Arlington Heights, Illinois, USA). The DNA templates were self-complementary hairpins (5' - TACCgTCgACgTCgACgCTggCg-AgCgTgCTgCggTTTT(C6-amino)TTACCgCAGCACgCTCgCCAgCg; SEQ ID NO:1). The reaction was performed in 100 µL volume at 37 °C with timepoints taken at 0, 1, 3, 5 and 10 min. The reaction products were electrophoresed down a denaturing (8 M urea) 20 % polyacrylamide gel and imaged on a typhoon phosphorimager. Complete single base extension was seen in 1 minute indicating efficient polymerase incorporation (indole linker gel, Fig. 6).

All patents, patent applications, and published references cited herein are hereby incorporated by reference in their entirety. While this invention has been particularly shown and described with references to preferred embodiments thereof, it will be understood by those skilled in the art that various changes in form and details may be made therein without departing from the scope of the invention encompassed by the appended claims.

25

- 33 -

CLAIMS

1. A nucleotide or nucleoside molecule, having a base that is linked to a detectable label via a cleavable linker.
5
2. The molecule of claim 1, wherein the base is a purine, or a pyrimidine.
- 10 3. The molecule of claim 1 or 2, wherein the base is a deazapurine.
4. The molecule of any of claims 1 to 3, having a ribose or deoxyribose sugar moiety.
15
5. The molecule of claim 4, wherein the ribose or deoxyribose sugar comprises a protecting group attached via the 2' or 3' oxygen atom.
- 20 6. The molecule of any of claims 1 to 5, that is a deoxyribonucleotide triphosphate.
7. The molecule of any of claims 1 to 6, wherein the detectable label is a fluorophore.
25
8. The molecule of any of claims 1 to 7, wherein the linker is acid labile, photolabile or contains a disulphide linkage.
- 30 9. A method of labeling a nucleic acid molecule, the method comprising incorporating into the nucleic acid molecule a nucleotide or nucleoside molecule, wherein the nucleotide or nucleoside molecule has a base that is linked to a detectable label via a

- 34 -

cleavable linker.

10. The method of claim 9, wherein said incorporating is accomplished via a terminal transferase, a polymerase or a reverse transcriptase.
11. The method of claim 9 or 10, wherein the base is a deazapurine.
12. The method of any of claims 9 to 11, wherein the nucleotide or nucleoside molecule has a ribose or deoxyribose sugar moiety, and wherein the ribose or deoxyribose sugar comprises a protecting group attached via the 2' or 3' oxygen atom and which can be modified or removed to expose a 3' OH group.
13. The method of any of claims 9 to 12, wherein the nucleotide is a deoxyribonucleotide triphosphate.
14. The method of any of claims 9 to 13, wherein the label is a fluorophore.
15. The method of any of claims 9 to 14, wherein the linker is acid labile, photolabile, or contains a disulphide linkage.
16. The method of any of claims 9 to 15, wherein the detectable label and/or the cleavable linker is of a size sufficient to prevent the incorporation of a second nucleotide or nucleoside into the nucleic acid molecule.

- 35 -

17. A method for determining the sequence of a target single-stranded polynucleotide, comprising monitoring the sequential incorporation of complementary nucleotides, wherein the nucleotides each have a base that is linked to a detectable label via a cleavable linker, and wherein the identity of each nucleotide incorporated is determined by detection of the label linked to the base, and subsequent removal of the label.
18. A method for determining the sequence of a target single-stranded polynucleotide, comprising:
- (a) providing nucleotides, wherein the nucleotides have a base that is linked to a detectable label via a cleavable linker, and wherein the detectable label linked to each type of nucleotide can be distinguished upon detection from the detectable label used for other types of nucleotides;
 - (b) incorporating a nucleotide into the complement of the target single stranded polynucleotide;
 - (c) detecting the label of the nucleotide of (b), thereby determining the type of nucleotide incorporated;
 - (d) removing the label of the nucleotide of (b); and
 - (e) optionally repeating steps (b) - (d) one or more times;
- thereby determining the sequence of a target single-stranded polynucleotide.

- 36 -

19. A method according to claim 17, wherein each of the nucleotides are brought into contact with the target sequentially, with removal of non-incorporated nucleotides prior to addition of the next nucleotide, and wherein detection and removal of the label is carried out either after addition of each nucleotide, or after addition of all four nucleotides.
20. A method according to claim 17, wherein each of the nucleotides are brought into contact with the target together simultaneously, and non-incorporated nucleotides are removed prior to detection and subsequent to removal of the label.
21. A method according to claim 17, comprising a first step and a second step, wherein in the first step, a first composition comprising two of the four nucleotides is brought into contact with the target, and non-incorporated nucleotides are removed prior to detection and subsequent to removal of the label, and wherein in the second step, a second composition comprising the two nucleotides not included in the first composition is brought into contact with the target, and non-incorporated nucleotides are removed prior to detection and subsequent to removal of the label, and wherein the first steps and the second step are optionally repeated one or more times.
22. A method according to claim 17, comprising a first step and a second step, wherein in the first step, a composition comprising one of the four nucleotides is brought into contact with the

- 37 -

target, and non-incorporated nucleotides are removed prior to detection and subsequent to removal of the label, and wherein in the second step, a second composition comprising the three
5 nucleotides not included in the first composition is brought into contact with the target, and non-incorporated nucleotides are removed prior to detection and subsequent to removal of the label, and wherein the first steps and the second step are
10 optionally repeated one or more times.

23. A method according to claim 17, comprising a first step and a second step, wherein in the first step, a first composition comprising three of the four
15 nucleotides is brought into contact with the target, and non-incorporated nucleotides are removed prior to detection and subsequent to removal of the label, and wherein in the second step, a composition comprising the nucleotide not included in the first composition is brought into
20 contact with the target, and non-incorporated nucleotides are removed prior to detection and subsequent to removal of the label, and wherein the first steps and the second step are optionally repeated one or more times.
25

24. A kit, comprising:


(a) individual nucleotides, wherein each
30 nucleotide has a base that is linked to a detectable label via a cleavable linker, and wherein the detectable label linked to each nucleotide can be distinguished upon detection from the detectable label used for

- 38 -


other three nucleotides; and
(b) packaging materials therefor.

- 5 25. The kit of claim 21, further comprising an enzyme
and buffers appropriate for the action of the
enzyme.


FIG. 1.




Uridine C5-linker




Cytidine C5-linker




N7 Deazaadenosine C7-linker



N7 Deazaguanosine C7-linker




Adenosine N6-linker



Cytidine N4-linker

where R_1 and R_2 , which may be the same or different, are each selected from H, OH, or any group which can be transformed into an OH.





X = H, phosphate, diphosphate or triphosphate





Guanosine N2-linker

FIG. 2.

Acid Labile Linkers




Disulfide Linker



X = O or N



R can be any substitution, including additional ring systems and systems in which the two marked R groups are linked together by further rings







the wavy bonds can symbolise anything
as they are not functionally important

FIG. 3.

Label ~~~ Cleavable linker ~~~ Base



Cleavable linkers may include:



R3 represents one or more substituents independently selected from alkyl, alkoxy, amino or halogen

Alternatively, cleavable linkers may be constructed from any labile functionality used on the 3'-block

where R_1 and R_2 , which may be the same or different, are each selected from H, OH, or any group which can be transformed into an OH, including a carbonyl

FIG. 4.

New disulfide cleavable linker (1)

Klenow exo- TMR
pH7.5 dUTP

+DTT



Incorporation Cleavage

AG AG

50mM Tris-HCl pH7.5, 10mM NaCl, 0.1mM EDTA
5mM MgCl₂, 2uM dNTP-fluor, 100nM SHP 5T hairpin AG oligo,
Klenow exo- (Amersham-Joyce) 10units.

t = 0, 30s, 1,3, 5

FIG. 3.

New Acid cleavable Sieber linker (3)

Klenow exo- TMR
pH7.5 dUTP

AG

50mM Tris-HCl pH7.5, 10mM NaCl, 2mM DTT, 0.1mM EDTA
5mM MgCl₂, 2uM dNTP-fluor, 100nM SHP 5T hairpin AG oligo,
Klenow exo- (Amersham-Joyce) 10units.

t = 0, 1, 3, 5, 10

FIG. 6.

New Acid cleavable Indole linker (4)

Klenow exo- TMR
pH7.5 dUTP



50mM Tris-HCl pH7.5, 10mM NaCl, 2mM DTT, 0.1mM EDTA
5mM MgCl₂, 2uM dNTP-fluor, 100nM SHP 5T hairpin AG oligo,
Klenow exo- (Amersham-Joyce) 10units.

t = 0, 1, 3, 5,